

Isotopic evidence for omnivory among European cave bears: Late Pleistocene *Ursus spelaeus* from the Peștera cu Oase, Romania

Michael P. Richards*[†], Martina Pacher[‡], Mathias Stiller[§], Jérôme Quilès[¶], Michael Hofreiter[§], Silviu Constantin^{||}, João Zilhão***, and Erik Trinkaus^{††‡‡}

Departments of *Human Evolution and [§]Evolutionary Genetics, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany; [†]Department of Archaeology, Durham University, South Park Road, Durham DH1 3LE, United Kingdom; [‡]Institut für Paläontologie, Universität Wien, Althanstrasse 14, 1090 Wien, Austria; [¶]Unité Mixte de Recherche 5608, Université Toulouse Le Mirail, 5 Allées Antonio Machado, 31058 Toulouse Cedex 9, France; ^{||}Institutul de Speologie "Emil Racoviță," Strasse Frumoas 31, 010986 Bucharest, Romania; ^{**}Department of Archaeology and Anthropology, University of Bristol, 43 Woodland Road, Bristol BS8 1UU, United Kingdom; and ^{††}Department of Anthropology, Campus Box 1114, Washington University, St. Louis, MO 63130

Contributed by Erik Trinkaus, November 21, 2007 (sent for review November 1, 2007)

Previous bone collagen carbon and nitrogen isotopic studies of Late Pleistocene European cave bears (*Ursus spelaeus*) have shown that these bears frequently had low nitrogen isotope values, similar to those of herbivores and indicating either unusual physiology related to hibernation or a herbivorous diet. Isotopic analysis of animal bone from the Peștera cu Oase (Cave with Bones), Romania, shows that most of its cave bears had higher nitrogen isotope values than the associated herbivores and were, therefore, omnivorous. The Oase bears are securely identified as cave bears by both their morphology and DNA sequences. Although many cave bear populations may have behaved like herbivores, the Oase isotopic data demonstrate that cave bears were capable of altering their diets to become omnivores or even carnivores. These data therefore broaden the dietary profile of *U. spelaeus* and raise questions about the nature of the carnivore guild in Pleistocene Europe.

carbon isotopes | nitrogen isotopes | diets | ancient DNA

Late Pleistocene cave bear (*Ursus spelaeus*) diet and physiology is not well known, although based on skull, mandible, and tooth morphology, they have been inferred to have been largely herbivorous (1–4). Moreover, isotopic studies of central and western European cave bears have shown that, in most cases where their values are compared with herbivores and carnivores from the same site, they have $\delta^{15}\text{N}$ values that plot with, or lower than, herbivores (Table 1). These dietary inferences are in the context of bears that had a period of dormancy, similar to modern black bears (*Ursus americanus*) and brown bears (*Ursus arctos*), given the abundance of hibernation nests in European caves. This pattern has been interpreted in several ways.

Based on comparisons of cave bear $\delta^{15}\text{N}$ values to herbivore values from three Late Pleistocene sites in France and to modern bear isotope data, Bocherens *et al.* (7) concluded that the cave bears were herbivores. They ruled out the possibility that hibernation had an effect on $\delta^{15}\text{N}$ values of modern bears, and by inference, it should not have affected cave bear $\delta^{15}\text{N}$ values.

Fernández *et al.* (12) analyzed $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ data from central and western European cave bears and hypothesized that the $\delta^{15}\text{N}$ values could be linked to climate. They found a link between $\delta^{15}\text{N}$ values and temperature, with cave bears from colder environments having lower $\delta^{15}\text{N}$ values. Although there are connections between climate and plant $\delta^{15}\text{N}$ values (17) that show that colder environments often have lower $\delta^{15}\text{N}$ values, they argued instead that the lower $\delta^{15}\text{N}$ values observed in Alpine cave bears were linked to hibernation. They argued that, because dormancy may result in lower cave bear $\delta^{15}\text{N}$ values, following Nelson *et al.* (11), colder environments require periods of longer dormancy that results in lower $\delta^{15}\text{N}$ values.

An alternative possibility (18) is that they had a dietary preference for nitrogen-fixing plants, which have lower $\delta^{15}\text{N}$ values. Such a diet might result in the very low $\delta^{15}\text{N}$ values seen in some cave bear samples (Table 1). However, there are dietary constraints in terms of potential protein intake, feeding rates, and digestibility of plant matter for at least modern brown bears (19–21), suggesting that a dependence on such plants does not fully explain the cave bear $\delta^{15}\text{N}$ pattern.

A fourth possibility, proposed by Hilderbrand *et al.* (14, 22) on the basis of higher $\delta^{15}\text{N}$ values from a diverse sample of cave bears (Table 1), is that some cave bears were largely herbivorous, as are some modern bears, but that other cave bears were more carnivorous and hence exhibit $\delta^{15}\text{N}$ values similar to those of many modern omnivorous *U. arctos* and *U. americanus*. In this interpretation, the variation in trophic level indicated by $\delta^{15}\text{N}$ values would reflect ecological plasticity among *U. spelaeus*, in which they were more or less carnivorous depending on available food resources, body-size requirements, interspecific competition, seasonality, and temperature.

Regardless of the cause, 86.1% of the available individual European adult cave bear $\delta^{15}\text{N}$ values ($n = 101$) are $<6.0\text{‰}$, and hence within the basically vegetarian range [the number rises to 95.6%, $n = 91$ without Hilderbrand *et al.*'s (14) 10 specimens] (Table 1). In the context of this distribution of $\delta^{15}\text{N}$ values, routine isotopic assessment of a cave bear metapodial associated with an early modern human cranium in the Late Pleistocene Peștera cu Oase, Caraș-Severin, southwestern Romania (23–27) [supporting information (SI) Text and SI Figs. 4 and 5], provided an anomalously high $\delta^{15}\text{N}$ value of 7.8‰ (SI Table 2, Oase O34-232). To further evaluate this result and address the cause of the generally low cave bear $\delta^{15}\text{N}$ values, we measured the isotope values of cave bears and associated fauna from the Peștera cu Oase.

Affinities of the Peștera cu Oase Ursid Remains. Because the primary concern of isotope analysis of Late Pleistocene *U. spelaeus* remains relates to the persistently low $\delta^{15}\text{N}$ values obtained for them, in contrast to the variable but generally higher (more

Author contributions: M.P.R., J.Z., and E.T. designed research; M.P.R., M.P., M.S., J.Q., M.H., S.C., J.Z., and E.T. performed research; M.P.R., M.S., J.Q., S.C., and E.T. analyzed data; and M.P.R., M.P., J.Z., and E.T. wrote the paper.

The authors declare no conflict of interest.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. EU289394–EU289402).

††To whom correspondence should be addressed. E-mail: trinkaus@artsci.wustl.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0711063105/DC1.

© 2008 by The National Academy of Sciences of the USA

Table 1. Previously determined isotope values of European cave bears, with the mean \pm SD for $N > 2$

Site	Region	$\delta^{13}\text{C}$	$\delta^{13}\text{C}$ range	$\delta^{15}\text{N}$	$\delta^{15}\text{N}$ range	N	Age, cal ka*	Ref
Liñares	NW Spain	-21.1 ± 0.3	-21.7 to -20.6	3.0 ± 0.4	2.1–3.6	20	40	5
Eirós	NW Spain	-21.1 ± 0.3	-21.3 to -20.4	5.1 ± 0.3	4.4–5.5	8	29	6
Aldène	S France	-20.8 ± 0.4	-21.3 to -20.5	3.1 ± 1.5	2.1–4.8	3	<30	7
Mialet	S France	-20.2 ± 0.4	-20.7 to -19.6	2.8 ± 1.6	0.8–4.4	5	—	7
Olaskoa	Pyrenees	-20.5	-20.5	2.7	2.7	1	12	7
Chauvet	S France	-20.5 ± 0.3	-21.2 to -20.1	3.8 ± 0.9	3.1–4.8	16	27.3–42.1	8
Scladina-1A	Belgium	-22.1 ± 0.2	-22.5 to -21.8	4.9 ± 1.2	3.0–6.1	7	44	9
Scladina-4	Belgium	-23.2 ± 0.3	-23.6 to -23.0	3.6 ± 0.6	3.0–4.5	4	120–130	10
Divje Babe	Slovenia	-20.5 ± 0.4	-19.9 to -21.0	1.9 ± 0.7	0.6–3.3	11	52	11
Ramesch	Austria	-22.5	-22.5	2.4	2.4	1	45–48	12
Herdengel	Austria	$-22.0, -22.1$	-22.1 to -22.0	2.3, 2.5	2.3–2.5	2	41.4–43.8	12
Schwabenreith	Austria	-22.9	-22.9	2.5	2.5	1	55–121	12
Lieglloch	Austria	$-20.1, -20.2$	-20.2 to -20.1	2.8, 4.1	2.8–4.1	2	30–40	12
Hartelsgraben	Austria	$-20.8, -21.0$	-21.0 to -20.8	0.8, 3.4	0.8–3.4	2	32–47	12
Coturines	Italy	-22.3 ± 0.9	-23.3 to -21.6	2.0 ± 0.5	1.6–2.5	3	45.3–51.3	12
Potočka Zijalka	Slovenia	-20.4	-20.4	4.2	4.2	1	35	12
Vindija	Croatia	$-20.7, -21.1$	-21.1 to -20.7	1.3, 1.5	1.3–1.5	2	>34	13
Muierii†	Romania	20.4 ± 0.2	-20.7 to -20.2	5.1 ± 1.6	3.4–7.3	4	44–53	
Cioclovina†	Romania	$-20.3, -20.4$	-20.3 to -20.4	3.1, 4.3	3.1–4.3	2	36, 45	
Various	Europe	-21.5 ± 0.5	-22.0 to -20.6	8.4 ± 0.5	7.7–9.1	10	—	14‡

All isotope data are from bone samples derived from mature individuals; data from teeth and immature individuals are not included here or in Fig. 3. The individual values are provided in SI Table 3.

*Radiocarbon ages were converted to calendar years by using CalPal (www.calpal.de).

†The Muierii and Cioclovina data are from OxA in the context of radiocarbon dating (15, 16).

‡The individual various European specimens are from Hilderbrand *et al.* (14) and were provided by Hilderbrand (personal communication).

omnivorous) values generally found for *U. arctos* (7, 28–30), it is essential to establish the taxonomic affinities of the bear bones from the Peștera cu Oase.

Morphological Affinities. A series of morphological contrasts between cave and brown bears are well established (31–34). These include, for *U. spelaeus*, broad molars with heavily wrinkled surfaces and additional cusps, loss of the first three premolars, pronounced doming of the forehead (especially in large males) (Fig. 1), large masseteric fossae and sagittal crests, large temporomandibular articulations, more robust long bones and metapodials, shorter extremities, pronounced tibial torsion, medial humeral curvature, and aspects of pedal facet morphology. All of these characteristics are found in all of the diagnostic bear remains from the Peștera cu Oase, and their morphology shows no distinctive differences from other cave bear samples that cannot be easily accommodated within models of normal, intraspecific temporal and spatial variation (25). There are no *U. arctos* in the sample.

Genetic Affinities. To provide additional confirmation of the morphological evidence, mitochondrial DNA (mtDNA) was extracted, amplified, and sequenced from 19 ursid samples (SI Table 2). All 19 individual sequences of the Peștera cu Oase ursids show clear affinity to central European cave bear sequences (35) rather than to brown bears. They do not form a monophyletic group within cave bear mtDNA variation, and the range of the Oase bear haplotypes is spread throughout most of the variability known for central European cave bear populations from southern Germany, Austria, Croatia, and Slovakia (35–37).

Stable Isotopes of the Peștera cu Oase *U. spelaeus* Remains. The Peștera cu Oase faunal isotope data are in Fig. 2 and SI Table 2. The herbivores (*Capra ibex* and *Cervus elaphus*) have values similar to those observed elsewhere for herbivores in Pleistocene Europe (38–40), with $\delta^{15}\text{N}$ values ranging between 2.2‰ and 5.4‰. The higher *Canis lupus* $\delta^{15}\text{N}$ values are similar to those for European carnivores (9, 10, 41, 42). Whereas the wolf and

deer remains come from the same time period as the bears, the ibex remains are more recent. There is fluctuation in faunal $\delta^{15}\text{N}$ values over the past 50,000 years (38), and it is not necessarily valid to compare fauna from different time periods. The variation observed in European fauna occurs principally at ≈ 13 thousand years before the present (ka BP) and in the Holocene, where herbivores have reduced $\delta^{15}\text{N}$ values; this may have affected the ibex scapula but probably not the other samples.

The Oase cave bear isotope data (Fig. 2 and SI Table 2) are shifted negatively in $\delta^{13}\text{C}$ values ($-21.5 \pm 0.4\text{‰}$, $n = 21$), but in this they match the values of most of the other European cave bear samples (Table 1 and SI Table 3). However, the Oase bear $\delta^{15}\text{N}$ values are principally between the relatively “carnivorous” values of 5.7–9.8‰, with one “herbivorous” value of 3.6‰ ($7.8 \pm 1.4\text{‰}$, $n = 21$; $8.0 \pm 1.1\text{‰}$, $n = 20$ without the one low value). A Wilcoxon nonparametric comparison of the pooled herbivore versus cave bear $\delta^{15}\text{N}$ values for the Oase sample provides a $P < 0.001$ ($P = 0.001$ if the more recent ibex scapula is deleted from the sample); $P = 0.036$ between the Oase bears and more carnivorous wolf $\delta^{15}\text{N}$ values. These cave bear isotopic values cannot be the product of diagenesis, because the other fauna, including the contemporaneous canid and cervid remains, have expected $\delta^{15}\text{N}$ values, and all have expected C/N values.

These Oase bear data contrast strongly with all previously published cases of cave bear $\delta^{15}\text{N}$ values associated with herbivore data from the same site, in which the cave bears have $\delta^{15}\text{N}$ values that are similar to, or lower than, herbivores from the same site. Most of the unpublished specimens from the Romanian sites of Cioclovina and Muierii (15–16) follow the same pattern (Table 1 and SI Table 3), although two specimens (Muierii 05-6 and 05-11) have $\delta^{15}\text{N}$ values of 6.0‰ and 7.3‰, and three specimens from Scladina 1A [$\delta^{15}\text{N}$ of 5.7‰, 6.0‰, and 6.1‰ (9)] are close (Fig. 3). The museum-derived cave bear sample analyzed by Hilderbrand *et al.* (14) produced $\delta^{15}\text{N}$ values similar to the Oase ones, although that may be due in part to selection for large (“museum quality”) specimens (22) because the degree of carnivory (and hence $\delta^{15}\text{N}$ values) tends to correlate with size, at least among North American brown bears (29).



Fig. 1. Left anterolateral views of Peștera cu Oase adult *U. spelaeus* crania. From top to bottom, N32.1 (female), O34.62b (female), N32.21 (male), and N37.49 (male). The protuberance on the N32.21 dorsal neurocranium is a stalagmite.

There is a modest inverse relationship between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in the Oase cave bears ($r = -0.708$, $P < 0.001$; without the one low outlier, $r = -0.554$, $P = 0.011$). This pattern is occasionally observed in isotopic studies, and it is often interpreted as a mixing line between two food sources (43). In this case, it would indicate that the cave bears had two distinct dietary protein sources, one being plant foods and the other being animals with low $\delta^{13}\text{C}$ and high $\delta^{15}\text{N}$ values. None of the Oase herbivore species could be the main source of animal protein for the cave bears, because the $\delta^{13}\text{C}$ values are too different (there is almost no shift in $\delta^{13}\text{C}$ value between prey and consumer). Whatever the source of the animal protein, it is apparent from the Oase cave bear isotope data that *U. spelaeus*, as with *U. arctos*, was not strictly vegetarian.




Fig. 2. Isotope data for cave bears and other fauna from the Peștera cu Oase, Romania. Filled diamonds, *U. spelaeus*; open squares, *C. lupus*; filled triangles, *C. elaphus*; open triangles, *C. ibex*. S-EVA values are used except for the specimens for which only OxA data are available.

Discussion

The Peștera cu Oase isotopic results indicate that cave bears do not always have low $\delta^{15}\text{N}$ values when compared with herbivores from the same site. Moreover, this pattern is evident, although less dominant, in samples from Muierii and Scladina 1A. Therefore, cave bears had the ability to consume animal protein and behaved like omnivores or even carnivores, even if in many cases they were herbivorous.

This inferred dietary flexibility for cave bears is evident in the stable isotope values within some samples, including Oase, Muierii, and Scladina 1A, as well as between samples. In this variation *U. spelaeus* closely parallels the range of isotopic values seen for modern European and North American brown bears (5, 7, 28–30) and the few data available for Late Pleistocene brown bears (10, 42).

As a consequence of these $\delta^{15}\text{N}$ values, the dietary ecology of modern, higher-latitude bears (excluding polar bears) is relevant for that of cave bears, especially the North American brown bears (*U. arctos*, including the Kodiak and grizzly bears) given their high-latitude range, body-size variation, occupation of regions with less human ecological impact than most of Eurasia, and extensive database. Brown bear diets range from almost completely vegetarian, including ones with substantial amounts of fruit/berries, to ones




Fig. 3. Distributions of $\delta^{15}\text{N}$ values for *U. spelaeus* remains from European OIS 4–2 Late Pleistocene sites (Table 1) and the Peștera cu Oase (SI Table 2). Samples: Liñares (1), Eirós (2), southern France and Pyrenees (3), Chauvet (4), Scladina 1A (5), Divje Babe (6), eastern Alps (7), Muierii and Cioclovina (8), European (9), and Oase (10).

containing a substantial amount of fish and/or ungulate meat (19–21, 29, 30, 44, 45). All aspects of their omnivorous diets have limitations in availability, potential feeding rates, and nutritional value in any given environment; adequate weight gain for survival, reproduction, and hibernation therefore depends on a mix of as many food resources as are available (19, 21). Meat consumption, in particular, varies widely among and within brown bear populations, due, among nonmaritime bears, to the availability of ungulate fauna (29, 30, 44, 45). Large adult males also appear to be more carnivorous than females or subadult bears (28, 29). North American black bears (*U. americanus*) appear to have similar plant/meat dietary proportions as brown bears (29), except that the larger brown bears are frequently more carnivorous when the prime meat is maritime (e.g., salmon) (46). This ecological flexibility of modern brown bears therefore makes an appropriate model to understand the range of isotopic values now available for European cave bears, both within and between site-specific samples.

The Oase bear isotope data also indicate that there is no need to invoke special species-wide effects, such as a dormancy effect or a reliance on specific plant foods, to explain the frequently low $\delta^{15}\text{N}$ for European Late Pleistocene cave bears.

These isotopic data also imply that *U. spelaeus* participated, at least occasionally, in the European Late Pleistocene carnivore guild. This would have brought it into competition with the purer carnivores, especially cave lions and hyenas, as well as the more omnivorous ones such as brown bears and wolves. A very large bear obtaining much of its protein from vertebrate prey would have a serious effect on the dynamics of that guild. Moreover, contemporaneous European humans, both late archaic and early modern, consistently exhibit high $\delta^{15}\text{N}$ values (13, 47, 48), and an additional large-bodied carnivore would have further intensified any human–carnivore interactions within the ecosystem.

Conclusions

Isotopic analysis of the Late Pleistocene cave bears from the Peștera cu Oase, Romania demonstrates that at least this bear population had omnivorous dietary adaptations. This is in contrast to most of the other published cave bear isotope data, which have been taken to show that they were herbivorous. The Oase data confirm that cave bears, as with modern brown bears, were omnivorous, although they may have frequently had dietary regimes that were primarily herbivorous. In an ecological context containing an abundance of large social carnivores, such as Late Pleistocene Europe, bears would have needed the dietary flexibility that is emerging from these stable isotope data. The *U. spelaeus* stable isotope data from the Peștera cu Oase, in conjunction with the data for cave bears elsewhere in Europe, now permit us to view them in a broader dietary framework and thereby raise questions as to their roles within the ecosystems of Pleistocene Europe.

Materials and Methods

The Peștera cu Oase Ursid Sample. The overall assessment of the Peștera cu Oase ursid sample is based principally on those remains excavated from the Panta Strâmoșilor, supplemented by surface materials in the other galleries of the Peștera cu Oase (see [SI Text](#), [SI Fig. 4](#) and [5](#)). For isotopic and ancient DNA analysis, 22 ursid, and 10 other large mammal remains were selected from both the excavated and surface remains (see [SI Table 2](#)). Fifteen of the bear bones derive from excavations in the Panta Strâmoșilor, whereas the others come from the surfaces of the adjacent galleries, some from hibernation nests. Other faunal species are rare within the Peștera cu Oase; sufficiently preserved bones for isotopic analysis include wolf ($n = 4$), ibex ($n = 4$), and red deer ($n = 2$). All of the specimens are bone and derive from fully mature individuals.

Given the mixed nature of the skeletal remains, there may be some duplication of individuals within species. The bear mtDNA haplotypes provide a minimum number of individuals of eight. The radiocarbon ages for the ibex provide a minimum of three individuals. The cervid hemimandibles, however, are right and left specimens, morphologically similar, may represent the same

individual, and have similar isotopic values and radiocarbon ages; they are nonetheless plotted separately.

Morphological Assessment. The faunal remains from the Peștera cu Oase were determined to skeletal element and species based on morphological characteristics. Apart from a few fragments, morphological and morphometric species identification was readily apparent for the >5,000 bear remains excavated in the Panta Strâmoșilor and collected from the cave surfaces.

Sample Collection. The analytical samples were derived from mature bone diaphyseal, rib, palatal, and basal mandibular cortical bone and removed by using a stainless-steel blade on a Dremel tool.

Ancient DNA. Ancient DNA was extracted from 170 to 750 mg of bone material as described in Hofreiter *et al.* (36) and Rohland and Hofreiter (49, 50). Two hundred eight-five base pairs (bp) of the mitochondrial control region were amplified by using regular PCR with primers and amplification conditions from Hofreiter *et al.* (35) or under multiplex PCR conditions as described by Rompler *et al.* (51) by using primers from Hofreiter *et al.* (35, 36). Amplification products were cloned by using the TOPO-TA cloning kit (Invitrogen) by following supplier recommendations. Four to eight clones of each amplification product were sequenced on an ABI 3730 by using the BigDye Terminator v1.1 Cycle Sequencing kit and M13 universal primers. Clone sequences were aligned by using BioEdit (52) and merged to reconstruct the complete 285-bp sequence for each individual. Every sequence position was determined from clone sequences derived from at least two primary amplification products to avoid errors caused by ancient DNA template damage (53, 54). In cases of consistent differences among clones derived from independent amplification products, a third primary amplification was used to clarify the correct state for each position.

Stable Isotopes. The key assumption behind the analysis of carbon and nitrogen stable isotopes is that bone collagen is formed throughout a mammal's life, and the carbon and nitrogen in collagen is ultimately derived from dietary protein (55–57). The isotope ratios of the two stable isotopes of carbon ($^{13}\text{C}/^{12}\text{C}$ or $\delta^{13}\text{C}$) and nitrogen ($^{15}\text{N}/^{14}\text{N}$ or $\delta^{15}\text{N}$) in dietary protein are commonly well preserved in European Late Pleistocene mammal bone collagen. Therefore, measurement of the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of mammal bone collagen can be used to determine the main sources of dietary protein over the lifetime of that mammal. The bone collagen isotope values likely reflect dietary protein consumed over the past 10–20 years of life for an adult long-lived mammal (58). Although used to assess whether the dietary protein carbon is derived from marine versus terrestrial sources or from C_3 or C_4 plants, it is the trophic level of the protein consumed that is of interest here. Because the $\delta^{15}\text{N}$ of bone collagen is $\approx 2\%$ to $\approx 4\%$ higher than dietary protein (59, 60), one can use it to determine the main protein sources in the diets of cave bears, especially compared with other fauna from the same site, to see whether dietary protein derived mainly from plants, as indicated by other isotopic studies of cave bears, or from higher trophic level protein sources. Of particular relevance here, experimental work with living bears (14) has shown that bone collagen $\delta^{15}\text{N}$ values are an accurate reflection of diet.

A potential source of error for paleontological samples is postdepositional collagen bacterial and/or chemical diagenesis. Bone samples were therefore pretreated to remove exogenous carbon and isolate the original collagen. The resulting collagen extract is measured for a number of collagen quality indicators, including collagen yield, C/N ratio, percent C, and percent N in the collagen extract (61, 62).

We extracted collagen from 26 mammalian bones from the Oase site using standard extraction procedures [i.e., modified Longin (63) method (64) with the addition of an ultrafiltration step (65)]. Isotope measurements were made by using a ThermoFinnigan elemental analyzer coupled to a Delta Plus XP isotope ratio mass spectrometer at the Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany. Carbon isotope ratio measurements ($\delta^{13}\text{C}$) were made relative to the vPDB standard, and the nitrogen isotope ratio measurements ($\delta^{15}\text{N}$) were made relative to the AIR standard. All samples were measured in duplicate, and the errors (1σ) are better than $\pm 0.2\%$.

Additional isotopic data produced at the Oxford Radiocarbon Accelerator Unit (ORAU) during sample pretreatment for radiocarbon dating were included for 12 specimens. The ORAU extraction procedures are similar to those that we used, including ultrafiltration [Brown *et al.* (65)]. All of the data reported here have C/N ratios that are within the accepted range of 2.9–3.6 (61).

In addition, 15 of the same specimens were AMS radiocarbon dated, principally by ORAU (OxA); two specimens were dated by the University of Vienna laboratory (VERA), one by using ultrafiltration and the other of an intact molar, which experience has shown provides results similar to bone dates run with ultrafiltration.

ACKNOWLEDGMENTS. Fieldwork at the Peștera cu Oase was undertaken through the Institutul de Speologie "Emil Racoviță," Academia Română; with the permission of the Direcția Monumente Istorice Si Muzeze, Ministerul Culturii si Cultelor (authorizations 181/2004 and 47/2005) and supported by National Science Foundation Grant BCS-0409194, Wenner-Gren Foundation Grant 7111, Washington University, the Leakey Foundation, the Instituto Português de Arqueologia, the Institut Royal des Sciences

Naturelles de Belgique, Romanian National Council for Academic Research Grant CNCIS 1258/2005), and Romanian Ministry for Education and Research Grant CEEC 627/2005). The fieldwork was made possible through the efforts of the Pro Acva Grup (Ș. Milota, M. Gherase, and L. Sarcină) and O. Moldovan. G. V. Hilderbrand provided their original cave bear isotopic data. The Max Planck Society funded the isotopic and aDNA research. To all we are grateful.

- Kurtén B (1976) *The Cave Bear Story* (Columbia Univ Press, New York).
- Mattson DJ (1998) Diet and morphology of extant and recently extinct northern bears. *Ursus* 10:479–496.
- Rabeder G, Nagel D, Pacher M (2000) *Der Höhlenbär* (Thorbecke, Stuttgart).
- Grandal A, López F (2005) Sexual dimorphism and autogenetic variation in the skull of the cave bear (*Ursus spelaeus* Rosenmüller) of the European Upper Pleistocene. *Geobios* 38:325–338.
- Vila M, Fernández D, López F, Grandal A, Vidal JR (1999) Paleoecological implications inferred from stable isotopic signatures ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) in bone collagen of *Ursus spelaeus* ROS.-HEIN. *Cab Lab Xeol Laxe* 24:73–87.
- Fernández D (1998) Biogeoquímica isotópica ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) del *Ursus Spelaeus* del yacimiento de Cova Eirós, Lugo. *Cab Lab Xeol Laxe* 23:237–249.
- Bocherens H, Fizet M, Mariotti A (1994) Diet, physiology and ecology of fossil mammals as inferred from stable carbon and nitrogen isotope biogeochemistry: Implications for Pleistocene bears. *Palaeogeog Palaeoclimatol Palaeoecol* 107:213–225.
- Bocherens H, Drucker DE, Billiou D, Geneste JM, van der Plicht J (2006) Bears and humans in Chauvet Cave (Vallon-Pont-d'Arc, Ardèche, France): Insights from stable isotopes and radiocarbon dating of bone collagen. *J Hum Evol* 50:370–376.
- Bocherens H, et al. (1997) Paleobiological implications of the isotopic signatures (^{13}C , ^{15}N) of the fossil mammal collagen in Scladina cave (Sclayn, Belgium). *Quatern Res* 48:370–380.
- Bocherens H, Billiou D, Mariotti A (1999) Palaeoenvironmental and palaeodietary implications of isotopic biogeochemistry of Last Interglacial Neanderthal and mammal bones in Scladina Cave (Belgium). *J Archaeol Sci* 26:599–607.
- Nelson DE, Angerbjörn A, Liden K, Turk I (1998) Stable isotopes and the metabolism of the European cave bear. *Oecologia* 116:177–181.
- Fernández D, Vila M, Grandal A (2001) Stable isotopes data (delta ^{13}C , delta ^{15}N) from the cave bear (*Ursus spelaeus*): A new approach to its palaeoenvironment and dormancy. *Proc R Soc Biol Sci* 268B:1159–1164.
- Richards MP, et al. (2000) Neanderthal diet at Vindija and Neanderthal predation: The evidence from stable isotopes. *Proc Natl Acad Sci USA* 97:7663–7666.
- Hilderbrand GV, et al. (1996) Use of stable isotopes to determine diets of living and extinct bears. *Can J Zool* 74:2080–2088.
- Soficaru A, Doboș A, Trinkaus E (2006) Early modern humans from the Peștera Muierii, Baia de Fier, Romania. *Proc Natl Acad Sci USA* 103:17196–17201.
- Soficaru A, Petrea C, Doboș A, Trinkaus E (2007) The human cranium from the Peștera Cioclovina Uscațã, Romania: Context, age, taphonomy, morphology and paleopathology. *Curr Anthropol* 48:611–619.
- Amundson R, et al. (2003) Global patterns of the isotopic composition of soil and plant nitrogen. *Global Biogeochem Cyc* 17:10311–10318.
- Vila M, Fernández D, Grandal A (2001) Cave bear's diet: A new hypothesis based on stable isotopes. *Cab Lab Xeol Laxe* 26:431–439.
- Rode KD, Robbins CT (2000) Why bears consume mixed diets during fruit abundance. *Can J Zool* 78:1640–1645.
- Welch CA, Keay J, Kendall KC, Robbins CT (1997) Constraints on frugivory by bears. *Ecology* 78:1105–1119.
- Rode KD, Robbins CT, Shipley LA (2001) Constraints on herbivory by grizzly bears. *Oecologia* 128:62–71.
- Hilderbrand GV, Robbins CT, Farley SD (1998) Response: Use of stable isotopes to determine diets of living and extinct bears. *Can J Zool* 76:2301–2303.
- Trinkaus E, et al. (2003) An early modern human from the Peștera cu Oase, Romania. *Proc Natl Acad Sci USA* 100:11231–11236.
- Trinkaus E, Milotaș, Rodrigo R, Gherase M, Moldovan O (2003) Early modern human cranial remains from the Peștera cu Oase, Romania. *J Hum Evol* 45:245–253.
- Quiñès, J et al. (2006) Cave bears (*Ursus spelaeus*) from the Peștera cu Oase (Banat, Romania): Paleobiology and taphonomy. *Comptes Rendus Palévol* 5:927–934.
- Rougier H, et al. (2007) Peștera cu Oase 2 and the cranial morphology of early modern Europeans. *Proc Natl Acad Sci USA* 104:1164–1170.
- Zilhão J, et al. (2007) in *Rethinking the Human Revolution*, eds Mellars P, Boyle K, Bar-Yosef O, Stringer C (McDonald Institute, Cambridge UK), pp 249–262.
- Hilderbrand GV, et al. (1999) The importance of meat, particularly salmon, to body size, population productivity, and conservation of North American brown bears. *Can J Zool* 77:132–138.
- Jacoby ME, et al. (1999) Trophic relations of brown and black bears in several western North American ecosystems. *J Wildlife Manag* 63:921–929.
- Mowat G, Heard DC (2006) Major components of grizzly bear diet across North America. *Can J Zool* 84:473–489.
- Altuna J (1973) Hallazgos de Oso Pardo (*Ursus arctos*, Mammalia) en cuevas del País Vasco. *Munibe* 25:121–157.
- Torres T, Quintero I, Gomez E, Mansilla H, Martinez C (1978) Estudio comparativo de las mandíbulas de *Ursus spelaeus*, Rosenmüller-Heinroth—*Ursus deningeri*, Von Reichenau y *Ursus arctos*, Linneo. *Bol Geol Min* 89-III:203–222.
- Torres T (1988) Osos (Mammalia, Carnivora, Ursidae) del Pleistoceno Ibérico (*U. deningeri* von Reichenau, *U. spelaeus* Rosenmüller Heinroth y *U. arctos* Linneo) *Bol Geol Min* 99:1–316.
- Chagneau J (1985) Contribution à l'étude des os des extrémités des pattes d'*Ursus deningeri romeiensis*. Comparaison avec *Ursus arctos* et *Ursus spelaeus*. *Bull Soc Anthropol Sud-Ouest* 20:1–85.
- Hofreiter M, et al. (2002) Ancient DNA analyses reveal high mitochondrial DNA sequence diversity and parallel morphological evolution of Late Pleistocene cave bears. *Mol Biol Evol* 19:1244–1250.
- Hofreiter M, et al. (2004) Evidence for reproductive isolation between cave bear populations. *Curr Biol* 14:40–43.
- Hofreiter M, et al. (2004) Lack of phylogeography in European mammals before the last glaciation. *Proc Natl Acad Sci USA* 101:12963–12968.
- Richards MP, Hedges REM (2003) Bone collagen d^{13}C and d^{15}N values of fauna from northwest Europe reflect palaeoclimatic variation over the last 40,000 years. *Palaeogeogr Palaeoclimatol Palaeoecol* 193:261–267.
- Drucker D, Bocherens H, Bridault A, Billiou D (2003) Carbon and nitrogen isotopic composition of red deer (*Cervus elaphus*) collagen as a tool for tracking palaeoenvironmental change during late glacial and early Holocene in northern Jura (France). *Palaeogeogr Palaeoclimatol Palaeoecol* 195:375–388.
- Hedges REM, Stevens RE, Richards MP (2004) Bone as a stable isotope archive for local climatic information. *Quatern Sci Rev* 23:959–965.
- Ambrose SH (1998) in *Neanderthals and Modern Humans in Western Asia*, eds Akazawa T, Aoki K, Bar-Yosef O (Plenum, New York), pp 277–289.
- Jacobi RM, Higham TFG, Bronk Ramsey C (2006) AMS radiocarbon dating of Middle and Upper Palaeolithic bone in the British Isles: Improved reliability by using ultrafiltration. *J Quatern Sci* 21:557–573.
- Szwarcz HP (1991) Some theoretical aspects of isotope paleodiet studies. *J Archaeol Sci* 18:261–275.
- Gau RJ, Case R, Penner DF, McLoughlin PD (2002) Feeding patterns of barren-ground grizzly bears in the central Canadian arctic. *Arctic* 56:339–344.
- MacHutchon AG, Wellwood DW (2003) Grizzly bear food habits in the northern Yukon, Canada. *Ursus* 14:225–235.
- Belant JL, Kielland K, Follmann EH, Adams LZ (2006) Interspecific resource partitioning in sympatric ursids. *Ecol Appl* 16:2333–2343.
- Richards MP, Pettitt PB, Stiner MC, Trinkaus E (2001) Stable isotope evidence for increasing dietary breadth in the European mid-Upper Paleolithic. *Proc Natl Acad Sci USA* 98:6528–6532.
- Beauval C, Lacrampe-Cuyaubère F, Maureille B, Trinkaus E (2006) Direct radiocarbon dating and stable isotopes of the Neanderthal femur from Les Rochers-de-Villeneuve (Lussac-les-Châteaux, Vienne). *Bull Mém Soc Anthropol Paris* 18:35–42.
- Rohland N, Hofreiter M (2007) Ancient DNA extraction from bones and teeth. *Nature Prot* 2:1756–1762.
- Rohland N, Hofreiter M (2007) Comparison and optimization of ancient DNA extraction. *Biotech* 42:343–352.
- Rompler H et al. (2006) Multiplex amplification of ancient DNA. *Nat Prot* 1:720–728.
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98.
- Hofreiter M, Jaenicke V, Serre D, Haeseler A, Pääbo S (2001) DNA sequences from multiple amplifications reveal artifacts induced by cytosine deamination in ancient DNA. *Nucleic Acids Res* 29:4793–4799.
- Stiller M, et al. (2006) Patterns of nucleotide misincorporations during enzymatic amplification and direct large-scale sequencing of ancient DNA. *Proc Natl Acad Sci USA* 103:13578–13584.
- Sealy J (2001) in *Handbook of Archaeological Sciences*, eds Brothwell DR, Pollard AM (Wiley, Chichester, UK), pp 269–279.
- Katzenberg MA (2000) in *The Biological Anthropology of the Human Skeleton*, eds Katzenberg MA, Saunders SR (Wiley, New York), pp 305–327.
- Ambrose SH, Norr L (1993) in *Prehistoric Human Bone: Archaeology at the Molecular Level*, eds Lambert J, Grupe G (Springer, New York), pp 1–37.
- Wild EM, et al. (2000) ^{14}C dating with the bomb peak: An application to forensic medicine. *Nucl Instr Methods Phys Res* 172B:944–950.
- Schoeninger M, DeNiro M (1984) Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochim Cosmochim Acta* 48:625–639.
- Szwarcz H, Schoeninger M (1991) Stable isotope analyses in human nutritional ecology. *Yrbk Phys Anthropol* 34:283–321.
- DeNiro MJ (1985) Post-mortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to paleodietary reconstruction. *Nature* 317:806–809.
- Ambrose SH (1990) Preparation and characterization of bone and tooth collagen for stable carbon and nitrogen isotope analysis. *J Archaeol Sci* 17:431–451.
- Longin R (1971) New method of collagen extraction for radiocarbon dating. *Nature* 230:241–242.
- Richards MP, Hedges REM (1999) Stable isotope evidence for similarities in the types of marine foods used by Late Mesolithic humans at sites along the Atlantic coast of Europe. *J Archaeol Sci* 26:717–722.
- Brown TA, Nelson DE, Vogel JS, Southon JR (1988) Improved collagen extraction by modified Longin method. *Radiocarbon* 30:171–177.